



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

February 16, 2017

Ms. Haley Sears
Regulatory Assistant
Control Solutions, Inc.
5903 Genoa-Red Bluff
Pasadena, TX 77507-1041

Subject: Notification per PRN 98-10 – Adding a Marketing Claim
Product Name: Prodiamine 0.58 Plus
EPA Registration Number: 53883-171
Application Date: July 20, 2016
Decision Number: 519573

Dear Ms. Sears:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 for the above referenced product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action requested falls within the scope of PRN 98-10.

The label submitted with the application has been stamped “Notification” and will be placed in our records.

Should you wish to add/retain a reference to the company’s website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA’s Office of Enforcement and Compliance.

If you have any questions, you may contact Kathryn Montague at 703-305-1243 or via email at montague.kathryn@epa.gov.

Sincerely,

A handwritten signature in black ink that reads "Kathryn V. Montague".

Kathryn Montague, Product Manager 23
Herbicide Branch
Registration Division (7505P)
Office of Pesticide Programs

NOTIFICATION

53883-171

The applicant has certified that no changes, other than those reported to the Agency have been made to the labeling. The Agency acknowledges this notification by letter dated:

02/16/2017

PRODIAMINE 0.58 PLUS

**FOR PRE-EMERGENCE CONTROL OF GRASS AND BROADLEAF WEEDS
IN LAWNS AND ORNAMENTAL TURF**

(excluding golf course putting greens)

**LANDSCAPE ORNAMENTALS, ESTABLISHED PERENNIALS AND
WILDFLOWER PLANTINGS**

**STOPS MANY WEED GRASSES BEFORE THEY GROW
PREVENTS CRABGRASS, ANNUAL BLUEGRASS, FOXTAIL**

ACTIVE INGREDIENT:

Prodiamine (CAS # 29091-21-2) 0.58%

INERT INGREDIENTS: 99.42%

TOTAL: 100.00%

KEEP OUT OF REACH OF CHILDREN

CAUTION

See back panel for further precautions and First Aid information

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Causes moderate eye injury (irritation). Harmful if inhaled or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Prolonged or repeated skin contact with product may cause allergic reactions in some individuals.

EPA REG NO. 53883-171

EPA EST. NO.

Control Solutions, Inc.
5903 Genoa-Red Bluff Rd.
Pasadena, TX 77507-1041
281-892-2500

NET CONTENT: _____

FIRST AID	
IF IN EYES:	<ul style="list-style-type: none"> ● Hold eye open and rinse slowly and gently with water for 15-20 minutes. ● Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. ● Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> ● Take off contaminated clothing. ● Rinse skin immediately with plenty of water for 15-20 minutes. ● Call a poison control center or doctor for treatment advice.
IF SWALLOWED:	<ul style="list-style-type: none"> ● Call a poison control center or physician immediately for treatment advice. ● Have person sip a glass of water if able to swallow. ● Do not induce vomiting unless told to do so by the poison control center or physician. ● Do not give anything by mouth to an unconscious person.
IF INHALED:	<ul style="list-style-type: none"> ● Move person to fresh air. ● If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. ● Call a poison control center or doctor for further treatment advice.
HOT LINE NUMBER	
<p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact SafetyCall® International (866) 897-8050 for emergency medical treatment information.</p>	

ENVIRONMENTAL HAZARDS

This product has low solubility in water. At the limits of solubility, this product is not toxic to fish. However, at concentrations above the level of water solubility, it may be toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters.

DIRECTION FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Not for use on plants being grown for sale or other commercial use, for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

This product is a pre-emergence herbicide that provides residual control of many grass and broadleaf weeds in established lawns and ornamental turf (excluding golf course putting greens),

landscape ornamentals and established perennials and wildflower plantings. It controls susceptible weeds by inhibiting weed seeds, germination and root development. When used according to directions, it will control; barnyardgrass; bluegrass; annual (*Poa annua*); carpetweed; chickweed, common; chickweed, mouseear (from seed); crabgrass (large, smooth); crowfootgrass; cupgrass, woolly; foxtails, annual; goosegrass¹; henbit; itchgrass; johnsongrass (from seed); junglerice; knotweed; kochia; lambsquarter, common; lovegrass; panicum (Texas, fall, browntop); pigweed; purslane, common; pusley, Florida; rescuegrass³, shepherd's purse², signalgrass, broadleaf; speedwell, Persian; sprangletop; spurge, prostrate; witchgrass; woodsorrel, yellow (from seed).


¹In many areas a single application of 2.5 to 6 pounds/1,000 square feet (112 to 259 pounds/acre) will control goosegrass. However, under heavy goosegrass pressure and/or an extended growing season, most effective weed control may be obtained with an initial application of 2.5 to 3.9 pounds/1,000 square feet (112 to 173 pounds/acre), followed after 60 to 90 days by a second application at doses that would not exceed the maximum annual rate (see "Minimum and Maximum Application Rate by Turfgrass Species" table in the Rates of Application section of this label).

²For this weed, apply in late summer, fall or winter prior to germination.

³Suppression only.

WHEN TO APPLY TO ESTABLISHED TURF: This product must be applied prior to germination of weed seeds, as it will not control established weeds.

THIS MAP SHOWS APPROXIMATE CRABGRASS GERMINATION DATES



APPLICATION DIRECTIONS: Apply this product uniformly, using a suitable spreader that has been properly calibrated. Most effective weed control will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation, or with shallow (1-2 inches) incorporation, prior to weed seed germination and within 14 days application.

USE PRECAUTIONS: The following precautions apply to the use of this product in turf grasses and lawns. Application of this product may thin emerged annual bluegrass and newly overseeded grasses. Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil (not thatch plus soil). Do not apply to newly set sod until the following year. Application to turf stressed by drought, low fertility, or pest damage may result in turf injury. Disturbing the herbicide barrier with cultural practices (such as core aeration) may result in reduced weed control. Do not apply to putting greens, or areas where dichondra, colonial bentgrass, velvet bentgrass or annual bluegrass (*Poa*

annua) are desirable species. Do not apply through any type of irrigation system. Do not apply aerially. Do not graze or feed livestock forage cut from treated areas.

RATES OF APPLICATION: This product is recommended for use on the turf grass species listed in the following table. The maximum amount of this product that may be applied per year is given for each turf grass species. Do not exceed the maximum yearly rate listed:

MINIMUM AND MAXIMUM APPLICATION RATES (by Turf Grass Species)		
Turf Species	Minimum Rate	Maximum Annual Rate¹
Bahiagrass, Bermudagrass ² Centipedegrass, Zoysiagrass, Seashore Paspalum, St. Augustinegrass Tall Fescue (including turf- Type)	4 lbs/1,000 sq. ft. (175 lbs/acre)	6 lbs/1,000 sq. ft. per year (259 lbs/acre per year)
Buffalograss Kentucky Bluegrass Perennial Ryegrass	2 lbs/1,000 sq. ft. (87.5 lbs/acre)	4 lbs/1,000 sq. ft. per year (173 lbs/acre per year)
Creeping Red Fescue	1.5 lbs/1,000 sq. ft. (65.3 lbs/acre)	2.9 lbs/1,000 sq. ft. per year (130 lbs/acre per year)
Creeping Bentgrass	1.5 lbs/1,000 sq. ft. (65.3 lbs/acre)	2.5 lbs/1,000 sq. ft. per year (112 lbs/acre per year)

¹These are the maximum rates per calendar year.

²May be used on newly sprigged or plugged Bermudagrass at rates not to exceed 2 pounds/1,000 square feet (87.5 pounds/acre). Newly sprigged or plugged Bermudagrass stolon rooting may be temporarily retarded. Suppression only of foxtail, goosegrass and rescuegrass due to reduced rates used in sprigging situations.

This product may be applied as a single application or in sequential applications to control weeds germinating throughout the year. Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rates recommended for each turf species. The length of time of residual weed control provided by this product is related to the rate applied. Use the higher rates listed for longer periods of weed control.

WHEN TO OVERSEED AFTER APPLICATION: This product will inhibit germination of turf seed if overseeded too soon after application. Follow rates and intervals in table for best overseeding/reseeding results. (See map above.)

USE RATE	MONTHS BEFORE OVERSEEDING		
	North	Transition	South
1.9 pounds/1,000 sq. ft. (86 pounds/acre)	4	4	4
2.5 pounds/1,000 sq. ft. (112 pounds/acre)	5	4	4
2.9 pounds/1,000 sq. ft. (130 pounds/acre)	6	5	5
3.16 pounds/1,000 sq. ft. (138 pounds/acre)	-	6	6
4 pounds/1,000 sq. ft. (173 pounds/acre)	-	7	7
4.5 pounds/1,000 sq. ft. (197 pounds/acre)	-	-	9
5.14 pounds/1,000 sq. ft. (224 pounds/acre)	-	-	10
5.94 pounds/1,000 sq. ft. (259 pounds/acre)	-	-	12

LANDSCAPE ORNAMENTALS: (including established perennials and wildflower plantings): This product may be applied for residual pre-emergence weed control in ornamentals. Apply at the rate of 2.5 to 5.94 pounds/1,000 square feet (112 to 259 pounds/acre) in fall and/or spring. Use higher rates of application for longer control periods. Sequential applications may be made

so long as the total amount of product applied does not exceed the maximum annual application rate of 5.94 pounds/1,000 square feet (259 pounds/acre) per year.

SPREADER SETTINGS to apply at the listed pounds per 1,000 square feet						
	1.9 lbs.	2.5 lbs.	3.16 lbs.	4 lbs.	5 lbs.	6 lbs.
Spreader A	X	X	X	X	X	X
(May contain additional spreader brands/models, spreader settings and spreader use instructions as appropriate for the product)						

APPLICATION TIMING AND INFORMATION: This product may be applied to newly transplanted and established ornamentals with a broadcast, over the top or directed application. Around new transplants, delay application to allow soil to settle and water thoroughly before applying. Apply after cuttings form roots and become established. Apply before budding/grafting or after buds/grafts have taken to avoid any inhibition of the tissue union. This product is a pre-emergence herbicide and will not control emerged weeds. Most effective weed control in ornamentals will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation, or with shallow (1 to 2 inch) mechanical incorporation, prior to weed seed germination and within 14 days following application. Best weed control is obtained when applied to soil free of clods, weeds and debris such as leaves. Prior to application, control existing vegetation by hand weeding, cultivation or the use of an appropriate post-emergence herbicide.

TOLERANT ORNAMENTAL SPECIES: This product will not harm most trees, shrubs, vines and flowers. The species listed below are tolerant to this product. Best results will be obtained when product is uniformly applied to the soil surface; avoid applying to ornamental shoots which may defeat the effort to get uniform distribution on the soil surface. This product may be applied over the top of these species. When plants are under stress (such as heat, drought or frost damage) some cultivars of listed plants may be sensitive to this product.

SCIENTIFIC NAME

Abies spp.
 Abelia grandiflora
 Acer palmatum
 A. Platanoides
 Achillea spp.
 Actindia chinensis
 Agapanthus africanus
 Agapanthus orientalis
 Akebiaq quintata
 Allium cemuum
 Anemone hybrida
 Aquilegia spp.
 Arctostaphylos densiflora
 Arctotheca calendula
 Artemesia spp.
 Aster spp.
 Aster X Frikartii
 Athrium Filiz-femina
 Aucuba japonica
 Begonia spp.
 Berberis gladwynensis
 B. Julianae
 B. mentorensis
 B. Thunbergil
 B. Verruculosa
 Bergenia cordifolia

COMMON NAME

Fir species** (Balsam, Frasier, Noble, etc.)
 Abelia: Sherwood
 Japanese Maple
 Norway Maple
 Yarrow: King Edward
 Kiwi*
 Lily-of-the-Nile (African Lily)
 Five-leaf or Chocolate Vine
 Lady's Leek, Nodding Onion
 Japanese Anemome
 Aquilegia: Red and Gold
 Vine Hill Manzanita
 Cape Weed
 Wormwood; Silver Mound, Castle
 Aster: Bonnie Blue, Purple Dome
 Lady Fern; Fern Lady
 Japanese Aucuba
 Fibrous Begonia: Hardy Grandis
 Barberry
 Wintergreen Barberry
 Mentor Barberry
 Japanese Barberry
 Warty Barberry

Boltonia asteroides
Buddleia davidii
Buxus microphylla
Callistemon viminalis
Calluna vulgaris
Campanula carpatica
Campis X Tagliabuana
Carpobrotus edulis
Cassia artemisoides
Ceanothus rigidus
Ceratostigma plumbagonoides
Chamaecyparis pisifera
Cleyera japonica
Citrus spp.
Coreopsis spp.
Cornus florida
C. Stolonifera
Cortaderia selloana
Cotoneaster apiculatus
C. buxifolius
C. dammeri
C. Microphyllus
Cretaegus spp.
Crococsmia spp.
Cupressus sempervirens
Delosperma alba
Delosperma spp.
Delphinium spp.
Dianthus deltooides
D. gratianopolitanus
Dodonea viscosa
Echinacea purpurea
Elaeagnus pungens
Euonymus fortunei
E. japonica
E. kiautschovica
Fatsia japonica
Forsythia intermedia
Forsythia suspensa
F. viridissima
Gaillardia spp.
Gardenia jasminoides
Gaura spp.
Gentiana dahurica
Geranium cinereum
Gladiolus spp.
Gysophyllia repens
Hedera helix
Hellanthenum spp.
Hemerocallis spp.
Hibiscus
Hibiscus spp.
Hibiscus rosa-sinensis
Hosta plantaginea
H. Sieboldiana
Houttuynia cordata var. variegata

Snowbank
Butterfly-Bush (Dwarf Blue); Royal Red
Japanese Boxwood
Weeping Bottlebrush
Scotch Heather
Tussock Bellflower, (White Clips)
Trumpet Flower, Madame Galen
Hottentot Fig (Ice Plant)
Feathery Cassia
Wild Lilac

False Cypress
Cleyera
Citrus species*
Coreopsis (Calliopsis) Early Sunrise, Moonbeam
Flowering Dogwood
American Dogwood
Pampas Grass
Cranberry Cotoneaster
Cotoneaster
Bearberry Cotoneaster
Rockspray Cotoneaster
Hawthorne
Lucifer
Italian Cypress
White Trailing Ice Plant
Cooperi Pink
Larkspur: Blue Elf
Dianthus: Maiden Pinks "Zing"
Cheddar Pink
Hop Bush
Coneflower, Purple; Magnus
Silverberry
Wintercreeper
Evergreen Euonymus
Spreading Euonymus
Japanese Aralia
Border Forsythia
Weeping Forsythia
Greenstem Forsythia
Gaillardia, Blanket Flower: "Goblin"
Gardenia, Cape-Jasmine

Gentian
Cranesbill
Gladiolus species**
Baby's Breath
English Ivy
Sunrose
Daylily: Aztec Gold, Stella de Oro, Tender Love
Rose of Sharon**
Mallow: Disco Belle White
Chinese Hibiscus
Hosta, Plantain Lily (Fragrant)
Hosta, "Seersucker"

Hydrangea macrophylla	Bigleaf Hydrangea
Ilex Bernyi	Holly
Ilex cornuta	Chinese Holly
I. crenata	Japanese Holly
I. opaca	American Holly
I. vomitoria	Yaupon Holly, Schillings
Inula ensifolia	
Iris ensata	Sword-Leaved Iris; Jodlesong
Iris siberica	Siberian Iris; Cabernet
Iris spp.	Iris species**
Jasminium nudiflorum	Winter Jasmine
Juniperus chinensis	Chinese Juniper; Nick's Compact, Parsonnii
J. conferta	Shore Juniper; Blue Pacific
J. davurica	Parsoni
J. horizontalis	Creeping Juniper
Juglans spp.	Walnut*
Justicia brandegeana	Shrimp Plant
Lagerstromia indica	Crape Myrtle
Lagerstromia indica X Fauriei	Crape Myrtle; Tuscarora
Lantana montevidensis	Weeping Lantana
Lavender spp.	Lavender; Munstead
Leontopodium alpinum	Edelweiss
Ligustrum amurense	Amur Privet
Ligustrum sinense	Chinese Privet; Variegata
L. japonicum	Japanese Privet
L. lucidum	Glossy Privet (Waxleaf)
Lilium spp.	Lily; Jazz
Liriope muscari	Big Blue Lily, Silver Mound, Evergreen Giant
Liriope muscari var. variegata	Liriope, Variegated
Liriope spicata	Liriope, Creeping
Lobelia Cardinalis	Cardinal Flower, Indian Pink
Lonicera japonica	Japanese Honeysuckle
L. tatarica	Tatarian Honeysuckle
Lorapetalum chinense	Burgundy
Lythrum spp.	Loosestrife; Modern Pink
Magnolia spp.	Magnolia species**
Maleophora luteola	Ice Plant
Malus spp.	Crabapple*
Miscanthus sinensis	Yaku Jima, Silberfeder
Nandina domestica	Heavenly Bamboo
Narcissus spp.	Narcissus species**
Nerium oleander	Oleander
Olea europaea	Olive*
Ophiopogon japonicus	Mondo Grass
Omanthus heterophyllus	Osmanthus (False Holly); Gulf Tide
Osteospermum fruticosum	Trailing African Daisy
Oxydendrum arboreum	Sourwood
Paeonia suffruticosa	Tree Peony
Pennisetum setaceum	Fountain Grass (Dwarf)
Perovskia atriplicifolia	
Persea americana	Avocado*
Photinia Fraseri	Frasier's Photinia (Redtip)
Physotegia Virginiana	Dragonhead, False; Vivid
Picea spp.	Spruce species** (Colorado Blue, Norway, etc.)
Pieris japonica	Lily-of-the-Valley Bush
Pinus brutia	Calabrian Pine

<i>P. canariensis</i>	Canary Island Pine
<i>P. Elliottii</i>	Slash Pine
<i>P. halepensis</i>	Aleppo Pine
<i>P. nigra</i>	Austrian Pine
<i>P. palustris</i>	Longleaf Pine
<i>P. radiata</i>	Monterey Pine
<i>P. strobus</i>	Eastern White Pine
<i>P. sylvestris</i>	Scotch Pine
<i>P. taeda</i>	Loblolly Pine
<i>P. Thunbergiana</i>	Japanese Black Pine
<i>P. Virginiana</i>	Virginia Pine
Pistachio spp.	Pistachio*
<i>Pittosporum rhombifolium</i>	Queensland Pittosporum
<i>P. tobira</i>	Japanese Pittosporum
<i>Podocarpus macrophyllus</i>	Japanese Yew
<i>Prunus laurocerasus</i>	English Laurel
<i>Prunus</i> spp.	Almond, Apricot, Nectarine, Peach, Plum and Prune*
<i>Pseudotsuga menziesii</i>	Douglas Fir**
<i>Pyracantha coccinea</i>	Firethorn (Scarlet)
<i>P. fortuneana</i>	Firethorn
<i>P. koidzumii</i>	Firethorn
<i>Pyrus</i> spp.	Bradford Pear spp.
<i>Quercus Shumardii</i>	Oak, Shumard's Red
<i>Quercus</i> spp.	Oak species
<i>Raphiolepis indica</i>	Indian Hawthorne
<i>Raphiolepis umbellata</i>	Yedda Hawthorne
<i>Rhododendron</i> (including Azalea)	'Coral Bells,' 'Formosa,' 'Hino-crimson,' 'PJM' 'Delaware Valley White,' 'Roseum Elegans,' 'Flame Creeper', 'Girard Crimson', 'Wakeiebisu' 'White Gumbo', 'George L. Tabor'
<i>Rosa banksiae</i>	Lady Bank's Rose
<i>Rosmarinus officinalis</i>	Rosemary*
<i>Rudbeckia</i> spp.	Black-Eyed Susan; Goldsturm
<i>Santolina virens</i>	Santolina
<i>Saxifraga</i> spp.	Saxifrage; Purple Dome
<i>Scabiosa</i> spp.	Pincushion Flower
<i>Sedum album</i>	Stonecrop
<i>Sedum caucicola</i>	Stonecrop; Lidakense
<i>Sedum dasyphyllum</i>	Stonecrop
<i>Sedum spurium</i>	Stonecrop; Dragon's Blood
<i>Spiraea bumalda</i>	Spirea; Anthony Waterer
<i>Syzygium paniculatum</i>	Australian Brushcherry
<i>Taxus cuspidata</i>	Japanese Yew
<i>T. media</i>	Yew
<i>Teucrium</i> spp.	Germander
<i>Thalictrum diptercarpum</i>	Meadow Rue
<i>Thuja occidentalis</i>	American Arborvitae
<i>Trachelospermum asiaticum</i>	Star Jasmine
<i>Tsuga canadensis</i>	Canada Hemlock
<i>Tulipa</i> spp.	Tulip species
<i>Veronica</i> spp.	Veronica, Speedwell; Sunny Border
<i>Viburnum japonicum</i>	Japanese Viburnum
<i>V. odoratissimum</i>	Sweet Viburnum
<i>V. plicatum</i>	Japanese Snowball
<i>V. rigidum</i>	Canary Island Viburnum

V. suspensum
V. tinus
V. trilobium
V. wrightii
Vinca major
Vinca Minor
Vitis spp.
Weigela florida
Yucca aloifolia
Y. filametosa

Arrowood Viburnum
Laurustinus
Cranberry Bush
Leatherleaf Viburnum
Greater Periwinkle, Vinca
Common Periwinkle, Vinca
Grape*
Old Fashioned Weigela
Spanish Bayonet, Yucca
Adam's Needle, Yucca

*Do not use on food producing trees, vines or plants.

**Not for use on container grown plants.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in original container in a locked storage area out of reach of children and domestic animals.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Non refillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Offer for recycling if available. If recycling is not available; then dispose of empty bag in a sanitary landfill, or if allowed by State and local authorities, by burning. If burned, say out of smoke.

CONDITIONS OF SALE AND WARRANTY

The DIRECTIONS FOR USE of this product are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop, injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of Control Solutions, Inc. or the Seller. All such risks shall be assumed by the Buyer.

Control Solutions, Inc. warrants that this product conforms to the chemical description on the label, and is reasonably fit for the purposes set forth in the DIRECTIONS FOR USE, when it is used in accordance with such directions, subject to the inherent risks mentioned above.

CONTROL SOLUTIONS, INC. NEITHER MAKES NOR INTENDS, NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE ANY OTHER WARRANTIES, EXPRESSED OR IMPLIED, AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. THIS WARRANTY EXTENDS TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS, WARNING, OR CAUTIONS. BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION TO REPLACEMENT OF, OR THE REPAYMENT OF THE PURCHASE PRICE FOR, THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INDIRECT OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. CONTROL SOLUTIONS,

INC. and the seller offer this product, and the Buyer and User accept it, subject to the foregoing
CONDITIONS OF SALE AND WARRANTY.

Optional Marketing Language: [Prevents Crabgrass]